ARCHIVES

OF

DERMATOLOGY AND SYPHILOLOGY

EDITORIAL BOARD

HOWARD FOX, Chief Editor
140 East Fifty-Fourth Street, New York 22

C. C. DENNIE, Kansas City, Mo.
HAROLD N. COLE, Cleveland
HERBERT RATTNER, Chicago
FRED D. WEIDMAN, Philadelphia
GEORGE M. MacKEE, New York
C. GUY LANE, Boston

VOLUME 50
1944

PUBLISHERS
AMERICAN MEDICAL ASSOCIATION
CHICAGO, ILL.

CONTENTS OF VOLUME 50

JULY 1944. NUMBER 1

PAGE

Penicillin Therapy of Impetigo Contagiosa and Allied Diseases: Use of Penicillium-Inoculated Dressing. Harold M. Johnson, M.D., Honolulu, Territory of Hawaii..... 1

Recurrent Vesicular Eruptions Appearing During Administration of Penicillin. Commander W. N. Graves (MC), U.S.N.R.; Lieutenant Commander C. C. Carpenter (MC), U.S.N.R., and Lieutenant (jg) R. W. Unangst (MC), U.S.N..... 6

Sulfur in Dermatologic Preparations. J. G. Downing, M.D.; L. M. Ohmart, A.M., and M. J. Stoklosa, B.S., Boston..... 8

Eczematous Contact Dermatitis Due to Mercurials: Report of a Case of a Reaction to Ten per Cent Ammoniated Mercury Ointment and Associated with Mercurial Poisoning. Captain M. H. Samitz, Medical Corps, Army of the United States..... 10

Immunization Therapy of Warts. Hans Biberstein, M.D., New York..... 12

Treatment of Psoriasis with Sarsaparilla Compound. Thomas S. Saunders, M.D., Portland, Ore. 23

An Unusual Form of Occupational Dermatitis: Report of an Outbreak in a Plant Manufacturing Hydrochloric Acid. Louis Schwartz, M.D., Bethesda, Md..... 25

Keratosis Follicularis Is Not Primarily a Follicular Disease. Francis A. Ellis, M.D., Baltimore 27

Syphilitic Hepatitis with Unusual Concomitant Manifestations: Report of Two Cases. Captain John N. Edson, Medical Corps, Army of the United States..... 31

Dermatitis from Lemon Grass Oil (Cymbopogon Citratus or Andropogon Citratus). H. Victor Mendelsohn, M.D., New York..... 34

Furuncular Myiasis. Oswaldo G. Costa, M.D., Bello Horizonte, Brazil..... 36

Triple Symptom Complex of Behcet: Report of a Case. Herman Ephraim, M.D., Haifa, Palestine 37

Dermatitis Due to Nail Polish: A Study of Twenty-Six Cases with the Chief Allergenic Component Tòluene Sulfonamide Formaldehyde Resin and Related Substances. Harry Keil, M.D., and Laird S. Van Dyck, M.D., New York..... 39

Clinical Notes:

Fixed Eruption from Sulfadiazine or Sulfamerazine. Hal E. Freeman, M.D., Long Beach, Calif..... 45

Abstracts from Current Literature..... 46

Society Transactions:

Cleveland Dermatological Society..... 50

Metropolitan Dermatological Society..... 52

Chicago Dermatological Society..... 55

Manhattan Dermatologic Society..... 67

San Francisco Dermatological Society..... 69

News and Comment..... 74

Book Reviews..... 74

Directory of Dermatologic Societies..... 76

AUGUST 1944. NUMBER 2

PAGE

Contact Testing of the Buccal Mucous Membrane for Stomatitis Venenata. Leon Goldman, M.D., Cincinnati, and Captain Bernard Goldman, Dental Corps, Army of the United States 79

Ichthyosiform Atrophy of the Skin in Hodgkin's Disease: Report of a Case, with Reference to Vitamin A Metabolism. A. J. Glazebrook, M.B., London, England, and W. Tomaszewski, M.D., Posen, Poland 85

Rat Mite Dermatitis: Acariasis Caused by the Tropical Rat Mite, Liponyssus Bacoti Hirst, 1914. C. Russell Anderson, M.D., Los Angeles 90

Tuberous Sclerosis: Report of a Case. Captain Zenas B. Noon and Captain Onie O. Williams, Medical Corps, Army of the United States 96

Kaposi's Varicelliform Eruption: Review of the Literature and Report of Two Cases of Its Occurrence in Adults. Robert L. Barton, M.D., and Louis A. Brunsting, M.D., Rochester, Minn. 99

Juxta-Articular Nodules. H. D. Chambers, M.D., Jamaica, British West Indies 105

Impetigo Herpetiformis in the Male: Report of One Case with Response to Sulfapyridine. A. Fletcher Hall, M.D., Santa Monica, Calif 107

Prurigo of Hebra (Severe Type). Ben Kanee, M.D., Vancouver, British Columbia, Canada 113

Superficial Epitheliomatosis: Report of a Case. Captain Stephen L. Weisman and Lieutenant Colonel Leon S. Medalia, Medical Corps, Army of the United States 117

Lichen Sclerosus et Atrophicus: Report of Its Occurrence in a Negro. S. Irgang, M.D., New York 120

Clinical Notes:
Contact Dermatitis Produced by Tincture of Merthiolate. Lester Hollander, M.D., Pittsburgh 123

Correspondence:
Coal Tar in Dermatologic Preparations. Herman Sharlit, M.D., New York 124

News and Comment 124

Abstracts from Current Literature 125

Society Transactions:
Baltimore-Washington Dermatological Society 128
New York Academy of Medicine, Section of Dermatology and Syphilis 133
New York Dermatological Society 138
Los Angeles Dermatological Society 145

Book Reviews 150

SEPTEMBER 1944. NUMBER 3

Erysipelothrix Rhusiopathiae Infection in Swine and in Human Beings: Comparative Study of the Cutaneous Lesions. Joseph V. Klauder, M.D., Philadelphia 151

Clinical Manifestations of Vitamin Deficiencies as Observed in the Federated Malay States. Paul Fasal, M.D., San Francisco 160

Epidermolysis Bullosa. Lieutenant Commander Justin L. Mooney, MC-V(S), U.S.N.R. .. 167

Dermatophytosis in Industry. Samuel M. Peck, M.D.; Isadore Botvinick, M.D., and Louis Schwartz, M.D., Bethesda, Md 170

Multiple Painful and Painless Glomus Tumors. Lieutenant Albert H. Slepyan, MC-V(S), U.S.N.R. 179

SEPTEMBER—*Continued*

PAGE

Connective Tissue Nevus. Karl Steiner, M.D., New York 183

Dyskeratosis Congenita with Pigmentation, Dystrophia Unguium and Leukoplakia Oris (Cole and Others): Report of the Cases of Two Brothers, with Improvement in the Leukoplakic Patches in One with Androgenic Medication. John Garb, M.D., and Gdali Rubin, M.D., New York 191

Clinical Notes:

Favorable Effects of Vitamin A in a Case of Acne of Long Duration. Thomas S. Saunders, M.D., Portland, Ore 199

Use of Sulfathiazole Suspension in the Isolation of Fungi. Leslie M. Smith, M.D., El Paso, Texas 199

Early Syphilis Masked and Delayed by Penicillin in the Treatment of Gonorrhea. Bertha Shafer, M.D., and Samuel J. Zakon, M.D., Chicago 200

Dermatitis Produced by Tintex Stocking Dye. Lester Hollander, M.D., Pittsburgh .. 200

Contact Dermatitis from Alfalfa and Bur Clover. C. Russell Anderson, M.D., Los Angeles 201

Pityriasis Rosea. Milton S. Hartman, M.D., Brooklyn 201

Correspondence:

Local Penicillin Therapy. Arthur G. Schoch, M.D., Dallas, Texas 202

Mycosis Fungoides. Jacob Tass, M.D., Jerusalem, Palestine 202

Abstracts from Current Literature 203

Society Transactions:

Cleveland Dermatological Society 206

Minnesota Dermatological Society 211

Brooklyn Dermatological Society 220

New York Academy of Medicine, Section of Dermatology and Syphilis 223

New York Dermatological Society 227

News and Comment 230

OCTOBER 1944. NUMBER 4

Cutaneous Leishmaniasis (Oriental Sore). D. A. Berberian, M.D., Beirut, Lebanon:

II. Incubation Period 231

III. Period of Infectivity of Saline Suspensions of Leishmania Tropica Cultures Kept at Room Temperature 233

IV. Vaccination Against Oriental Sore with Suspensions of Killed Leishmania Tropica 234

Cutaneous Leishmaniasis on the Isthmus of Panama. Captain B. H. Kean, Medical Corps, Army of the United States 237

Vaccination Against Oriental Sore: Report of Results of Five Hundred and Fifty-Five Inoculations. I. Katzenellenbogen, M.D., Jerusalem, Palestine 239

Treatment of Trichophytosis with Ethyl Chloride. Major J. H. Lewis and Captain William J. Morginson, Medical Corps, Army of the United States 243

Dermatitis from Tyroglyphidae in Handlers of Straw. Thomas S. Saunders, M.D., Portland, Ore. 245

Penicillin in Treatment of Simultaneous Infections of Syphilis and Gonorrhea. Captain Orlando Canizares, Medical Corps, Army of the United States 246

Cholinogenic Urticaria and Cholinogenic Itching. Ruben Nomland, M.D., Iowa City 247

OCTOBER—*Continued*

PAGE

Local Treatment of Psoriasis by Choline Chloride: Report of Two Cases. Harry H. Fox, M.D., Cincinnati 250

Treatment of Eczematous Contact Dermatitis with Intravenous Injections of Sodium Thiosulfate. Albert Strickler, M.D., Philadelphia 251

Scleropoikiloderma with Calcinosis Cutis, Raynaud-Like Syndrome and Atrophoderma. Ben Kanee, M.D., Vancouver, British Columbia, Canada 254

Effect of Folliculoid Hormones on Abnormal Skin: Further Observations of the Effect of Estradiol on the Skin of Mice of the Rhino, Hairless and Naked Strains. Hans Selye, M.D., Ph.D., D.Sc., F.R.S.(C), Montreal, Canada 261

Preparation of Material for Laboratory Diagnosis of Some Tropical Diseases of the Skin. Leon Goldman, M.D., Cincinnati 264

Dystrophia Unguium Mediana Canaliformis: Report of a Case. Lester P. Fowle, M.D., Lewisburg, Pa., and Lieutenant Richter H. Wiggall (MC), U.S.N. 267

Clinical Notes:

Rosacea in Twins. Thomas S. Saunders, M.D., Portland, Ore 269

Suppression of Syphilis by Penicillin Therapy of Gonorrhea. Lieutenant Commander Hugh E. Hailey (MC), U.S.N.R 269

Obituaries:

Frederick H. Dillingham, M.D 270

News and Comment 270

Correspondence:

Contact Eczema Due to Nail Polish. William L. Dobes, M.D., Atlanta, Ga., and Philip H. Nippert, Lieutenant, MC-V(S), U.S.N.R 270

Abstracts from Current Literature 271

Society Transactions:

New York Academy of Medicine, Section of Dermatology and Syphilis 274

New York Dermatological Society 283

Book Reviews 288

NOVEMBER 1944. NUMBER 5

Hazards of the External Use of Sulfonamide Compounds. E. William Abramowitz, M.D., New York 289

Partial Albinism and Nystagmus in Negroes: Report of Two Cases. Lieutenant Colonel L. J. A. Loewenthal, South African Medical Corps 300

Agranulocytosis Following Mapharsen Therapy: Report of Two Cases. Lieutenant Colonel Milosh Kasich, Medical Corps, Army of the United States 302

Poison Ivy Smoke: Experiments Demonstrating That Poison Ivy Smoke Is Not a Cause of Clinical Ivy Dermatitis. J. B. Howell, M.D., Dallas, Texas 306

John Gorrie, M.D., and an Early Case of Keloid. Irving S. Cutter, M.D., Chicago 308

Effect of Grenz Rays on Leprous Infiltrations: Report of an Attempt to Influence Leprous Infiltrations by Roentgen Rays of Long Wavelength. Felix Sagher, M.D., Jerusalem, Palestine 311

Spectrographic Analysis of Gold in the Skin Following Treatment of Lupus Erythematosus. Lawrence G. Beinhauer, M.D.; Frederick M. Jacob, M.D., and Palmer L. Beebe, M.D., Pittsburgh 315

NOVEMBER—*Continued*

PAGE

Aplastic Anemia with Acute Agranulocytosis, Thrombopenic Purpura and Complicating Mapharsen Therapy: Report of a Case, with Pathologic Observations. Hal E. Freeman, M.D., Long Beach, Calif. 320

Granuloma Annulare and Measles. Captain Norman M. O'Farrell, Medical Corps, Army of the United States 323

Unsuccessful Treatment of American Leishmaniasis with Penicillin: Report of a Case. Major James S. Snow, Medical Corps, Army of the United States 324

Dermatitis from Penicillin: Report of Two Cases. George W. Binkley, M.D., and Arnold Brockmole, M.D., Cleveland 326

Clinical Notes:

Mutilating Form of American Leishmaniasis. Oswaldo G. Costa, M.D., Bello Horizonte, Brazil 328

Contact Dermatitis from Amorphous Sodium Penicillin. Seymour H. Silvers, M.D., Brooklyn 328

Correspondence:

Scleroderma with Bullae. H. J. Templeton, M.D., Oakland, Calif. 330

Local Penicillin Therapy. Adolph Rostenberg Jr., M.D., Washington, D. C. 330

Rat Mite Dermatitis. C. Russell Anderson, M.D., Los Angeles 330

News and Comment 330

Obituaries:

Jerome Kingsbury, M.D. 331

Andrew Porter Biddle, M.D. 333

Abstracts from Current Literature 334

Society Transactions:

Cleveland Dermatological Society 337

Philadelphia Dermatological Society 342

Book Reviews 354

DECEMBER 1944. NUMBER 6

Exit Dose in Dermatologic Roentgen Therapy. George C. Andrews, M.D.; Carl B. Braestrup, B.S., P.E., and Eldred B. Heisel, M.D., New York 355

Combined Bodian and Masson Staining Method Applied to Skin. William B. Dublin, M.D., Los Angeles 361

Erythema Elevatum Diutinum. Lloyd W. Ketron, M.D., Baltimore 363

Nevus Anemicus (Voerner). F. O. Piorkowski, M.D., L.R.C.P., L.R.C.S. (Edinburgh), Nairobi, Kenya 374

Postwar Graduate Training in Dermatology and Syphilology. C. Guy Lane, M.D., Boston, and George M. Lewis, M.D., New York 378

American Onchocerciasis. Leon Goldman, M.D., Cincinnati 385

American Leishmaniasis of Frambesial Type. Oswaldo G. Costa, M.D., Bello Horizonte, Brazil 394

CIII. Kaposi's Varicelliform Eruption: Report of Five Cases. Clinton W. Lane, M.D., and Walter C. Herold, M.D., St. Louis 396

DECEMBER—*Continued*

PAGE

Ptosis of a Kidney as a Cause of Pruritus Vulvae. Albert Strickler, M.D., Philadelphia.. 405

Dermatologists of the United States and Canada in Active Service in the Present War: Supplementary List.... 406

Obituaries:
- John Borneman Ludy, M.D.... 410
- Laird Sumner Van Dyck, M.D.... 411

News and Comment:.... 412

Abstracts from Current Literature.... 413

Society Transactions:
- Cleveland Dermatological Society.... 416
- Metropolitan Dermatological Society.... 421
- Chicago Dermatological Society.... 422
- Minnesota Dermatological Society.... 429
- New England Dermatological Society.... 433
- Los Angeles Dermatological Society.... 439

Book Reviews.... 442

General Index.... 443

A Radio Program of Interest to All Physicians . . .

"THE DOCTOR FIGHTS"

starring RAYMOND MASSEY

HERE is a Report to the Nation on the wide-spread activities of America's doctors in a world at war, not only on the battlefronts, but on the home front as well. Documentary histories of medical heroism, carefully authenticated and ethically presented, should prove of interest to every physician, military or civilian. The comments or suggestions of the profession are welcomed.

Tuesday Evenings
COLUMBIA BROADCASTING SYSTEM
9:30 E.W.T.—8:30 C.W.T.
7:30 M.W.T.—6:30 P.W.T.

SCHENLEY LABORATORIES, INC. · *Producers of* **PENICILLIN** *Schenley*
Lawrenceburg, Ind.

B R E C K



BRECK WATER RESISTANT CREAM

Water Resistant Cream forms a film of material on the hands which is resistant to the action of water and water soluble substances All that is necessary is to rub the cream on the hands and then water appears to run off the hands like water off a duck's back When men or women must work with their hands in water or must handle damp or wet material Breck Water Resistant Cream will help keep hands in good condition · This comparatively new way of giving protection to the hands of workers in War Industrial Plants we like to believe is helping workers and management alike in the war effort

JOHN H BRECK INC · MANUFACTURING CHEMISTS · SPRINGFIELD 3 MASSACHUSETTS
CANADIAN ADDRESS · 47 CLARENCE STREET · OTTAWA

More dermatologists buy
Mennen Brushless Shave
for their personal use
than any other brand,
more than the next two
leading brands combined.

Results of a survey among the
Dermatologists on the official
A. M. A. roster

MENNEN

An Achievement in
PROTECTION AGAINST CONTAMINATION

1—Instead of the two-liter flasks in which penicillin ordinarily is made by "surface culture," Peniciliin-C.S.C. is made in a battery of giant tanks, each of 12,000 gallon capacity, by "submerged culture," an operation of vastly increased sensitivity, calling for the utmost in care and control. **2**—Vial-filling; note the safeguards against contamination. **3**—Cold room, where Penicillin-C.S.C. is frozen prior to vacuum-drying.



1



2

3



4

5

4—The "last word" in controlled vacuum-drying equipment. The number of these evaporators indicates the magnitude of Penicillin-C.S.C. production. **5**—Vial-sealing and capping.

THE "Sterile Area" IN THE PENICILLIN · C.S.C. PLANT

WALLS of highly polished opal glass and translucent glass brick, and rounded floor and ceiling abutments, permitting of maximum cleanliness—

air-conditioning that controls temperature, humidity, and particle content—

sterilizing lamps that destroy air-borne microorganisms—

sterilizing-lamp-controlled "locks" that prevent undue airflow from room to room—

sterile clothing (masks, gowns, shoes, gloves) worn by all technicians—

facial shields which carry the technician's breath away from the work area—

these are but a partial list of the safeguards employed in the "sterile area" of the C.S.C. plant.

Out of its quarter-century of research and experience in microbiotic production, Commercial Solvents Corporation has developed not only these safeguards, but also the "submerged culture" method which produces Penicillin-C.S.C. in giant three-story tanks.

This combination of mass production methods, skilled personnel, the utmost in safeguards, and unremitting laboratory control spells two assurances—

Penicillin-C.S.C. will always be of dependable potency, sterility, and pyrogen-freedom—

Penicillin-C. S. C., now allocated as the armed forces direct, will be available in adequate distribution throughout the country as soon as released.

Testing for potency

Testing for toxicity



Testing for pyrogens

PHARMACEUTICAL DIVISION

COMMERCIAL SOLVENTS *Corporation*

Penicillin Plant
Terre Haute, Ind.

CSC

17 East 42nd Street
New York 17, N. Y.



Capacity conservatively rated at 40,000,000,000 (forty billion) Oxford Units per month.

Accepted by the Council on Pharmacy and Chemistry



For the Treatment of Neurosyphilis

Aldarsone is an organic compound of pentavalent arsenic having pronounced spirocheticidal and trypanocidal properties. Administered intravenously, either alone or combined with artificial fever therapy,* the drug is indicated for the treatment of central nervous system syphilis. Aldarsone is supplied in 0.5-Gm. and 1.0-Gm. ampoules. It is soluble in sterile distilled water and does not require neutralization with alkali. Complete literature will be sent on request. ABBOTT LABORATORIES, North Chicago, Ill.

**Venereal Disease Information, 25:69, March, 1944.*

Aldarsone

REG. U. S. PAT. OFF.

(Phenarsone Sulfoxylate, Abbott)

ECZEMA

. . . AND THE COSMETIC FACTOR

Identifying the causative agent in cases of eczema may be extremely difficult. Many physicians have found it a wise precaution to safeguard their treatment regimen by prescribing Marcelle hypo-allergenic Cosmetics, since cosmetic allergens may be at fault. Known allergens have been removed or reduced to a minimum in the production of Marcelle hypo-allergenic Cosmetics.

Physicians have been prescribing Marcelle hypo-allergenic Cosmetics for years as a routine measure in allergic and suspected allergic cases. The importance of the cosmetic facter is evidenced by the success which physicians have experienced through the recommendation of Marcelle hypo-allergenic Cosmetics.

Marcelle hypo-allergenic Cosmetics have been accepted for advertising in publications of the American Medical Association for 12 years.

Write for professional samples and a formulary booklet specifying ingredients of Marcelle Cosmetics. This offer is not good in Canada.

MARCELLE COSMETICS, Inc.
1741 North Western Ave., Chicago 47, Illinois

Marcelle
HYPO-ALLERGENIC
COSMETICS

SULFATHIAZOLE *for Gonorrhea*

- **Cure rate better than 80 per cent.**
- **Total dose 20 grams, given over 5- or 10-day period.**
- **Amazingly low cost for medication.**

ALTHOUGH other sulfonamides are used, sulfathiazole is generally preferred for treatment of acute and chronic gonorrhea. Given orally, as directed, it has a high cure rate, most patients can be kept ambulant, and the cost of medication is amazingly low.

Sulfathiazole has made it possible for any practitioner or health officer, however remote from the laboratory, to control gonorrhea.

An ample quantity of Sulfathiazole Squibb is now available for hospitals and clinics wherever located. Manufactured in the Squibb Laboratories and hence subject to the rigid Squibb system of control, its purity, uniformity and efficacy are assured.

WHENEVER YOU
THINK OF SULFONAMIDES
THINK OF SQUIBB
Specify Sulfathiazole Squibb
Supplied in 0.5-Gm. tablets in bottles of 100 and 1000.

E·R·SQUIBB & SONS

Manufacturing Chemists to the Medical Profession since 1858

Luzier Cosmetics and Allergy

Women use cosmetics because they have developed a need for them: they are essential to modern standards of good-grooming and therefore contribute to a sense of well-being. Your patient's appearance, viewed cosmetically, is a factor that deserves your consideration both during hospitalization and convalescence. Cosmetics cannot lift faces, but they certainly perform wonders when it comes to lifting a woman's spirits. Women have an instinctive desire to look pretty and to smell sweet.

Since cosmetics are so universally used it is not to be wondered that they sometimes figure in the field of allergy. That is why when there is a history of allergy we suggest that patch tests be made with those of our products the subject is using or contemplates using. If they test positive, further testing with their constituents is indicated to determine the offending agents. These found, we frequently can modify our formulas to suit the subject's requirements. The patch test is generally considered best for testing cosmetics because it most closely approximates the conditions under which they are normally used.

While our products are free from so-called common cosmetic allergens, such as orris root and rice starch, we feel it should be made clear that any of their normally innocuous ingredients might be allergenic to the allergic individual. It is our practice to write our patrons a letter to this effect when a history of allergy is involved.

It is our experience that many persons with allergic constitutions cannot tolerate scented cosmetics; therefore we routinely recommend and select unscented products when there is a history or suspicion of allergy. This practice is not to imply or suggest that the subject is sensitized to perfume; it is solely to safeguard against the possibility.

In specific cases of allergy or suspected allergy, when the subject is using or contemplates using our products, we are pleased on his request to send her doctor the involved raw materials for patch testing, also such information concerning our products as may have a bearing on the case.

Since in the light of present knowledge it is not possible, save in specific cases, to make non-allergenic cosmetics, we believe the cosmetic requirements of the allergic individual should be considered by her doctor in the light of the formulas and general characteristics of the products she is using or contemplates using.

Luzier's, Inc., Makers of Fine Cosmetics & Perfumes

KANSAS CITY, MISSOURI

For Surgical Antisepsis

Zephiran Chloride is a germicide of high bactericidal and bacteriostatic potency. In proper dilutions it is nonirritating and relatively nontoxic to tissue cells.

Zephiran Chloride possesses detergent, keratolytic and emulsifying properties, which favor penetration of tissue surfaces, hence removing dirt, skin fats and desquamating skin.

INDICATIONS

Zephiran Chloride is widely employed for skin and mucous membrane antisepsis—for preoperative disinfection of skin, denuded skin and mucous membranes, for vaginal instillation and irrigation, for vesical and urethral irrigation, for wet dressings, for irrigation in eye, ear, nose and throat infections, etc.

HOW SUPPLIED

Zephiran Chloride is available in

TINCTURE 1:1000 Tinted

TINCTURE 1:1000 Stainless

AQUEOUS SOLUTION 1:1000

in 8 ounce and 1 gallon bottles.

Write for informative booklet

WINTHROP CHEMICAL COMPANY, INC.

Pharmaceuticals of merit for the physician

NEW YORK 13, N. Y. • WINDSOR, ONT.

ZEPHIRAN CHLORIDE

"Zephiran" Trademark Reg. U.S. Pat. Off. & Canada

Brand of BENZALKONIUM CHLORIDE REFINED